



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$7=1+6=2+5=3+4=4+3=5+2=6+1$; hence the chance against throwing 7 at one throw is $1-\frac{6}{36}=\frac{5}{6}$.

Hence, the chance against throwing either 7 or 11 is $1-(\frac{1}{18}+\frac{1}{6})=\frac{7}{9}$.

II. Solution by J. SCHEFFER, A. M., Hagerstown, Md.

The chance of throwing 7 with two dice is $\frac{6}{36}=\frac{1}{6}$, and that of throwing 11 is $\frac{2}{36}=\frac{1}{18}$; therefore, the chance of throwing either 7 or 11 is $\frac{1}{6}+\frac{1}{18}=\frac{2}{9}$.

\therefore the odds against this event $=7:2$.

III. Solution by WALTER H. DRANE, Graduate Student, Harvard University, Cambridge, Mass.

The combinations which give 7 or 11 with two dice are four in number, viz.: $6+1$, $5+2$, $4+3$, and $6+5$.

The total number of combinations is 36. Hence the chance of throwing either a 7 or 11 at one throw is $\frac{4}{36}=\frac{1}{9}$. Hence the odds are 8 to 1 against the event.

IV. Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

7 can be thrown 6 ways at one throw with two dice, as follows:

6 and 1, 1 and 6, 5 and 2, 2 and 5, 4 and 3, 3 and 4.

\therefore chance of throwing 7 is $\frac{6}{36}=\frac{1}{6}$.

\therefore the odds are 5 to 1 against the event.

11 can be thrown 2 ways, as follows: 6 and 5, and 5 and 6.

\therefore chance of throwing 11 is $\frac{2}{36}=\frac{1}{18}$.

\therefore the odds are 17 to 1 against the event.

7 is the most likely throw of all at one time with two dice.

V. Solution by ELMER SCHUYLER, Annapolis, Md.

$7=1+6=2+5=3+4$, each occurring in two ways.

\therefore 7 can occur in six ways.

$11=5+6$, can occur in 2 ways.

\therefore 7 and 11 can occur in eight ways.

$\therefore \frac{8}{36}=\frac{2}{9}$ = probability in favor of the event.

\therefore the odds are 7 to 2.

69. Proposed by Rev. W. ALLEN WHITWORTH, M. A.

There are n equal sugar sticks. Each stick is broken into two pieces, all positions of the fracture being equally likely. Of the two n pieces thus formed, a child is to take the largest. Show that his expectation is $[2n+1]/[2(n+1)]$ of a stick. [From *The Educational Times*, June, 1898.]

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

Let each stick be of unit length. Then since there are $2n$ pieces, and each piece can have any length from zero to unity, we have for the required expectation: